

How can the effects of toxics be prevented and treated?

It is possible to intervene at several levels to rid the body of toxic substances or their conversion products, including those involved in its physiological functions.

How can toxics be cleared from the body?



The “medicine cabinet” of an occupational health and safety department for nuclear operators – here at the CEA Saclay Centre – comprises mainly potassium iodide, Prussian blue and different forms of DTPA.

Cyrille Dupont/CEA

Any **exogenous** substance that enters a living organism is transferred and converted (**metabolised**), bound for a certain time, and eliminated at some variable rate either in its initial form or as a metabolite.

If this substance or its **metabolites** are not involved in normal physiological cell functions, they can induce an immediate or delayed pathological disorder: the substance is referred to as a toxin (from the Greek *toxikon*: poison).

Some **elements** are metabolised in a similar way to those the body normally uses. Strontium-90 (^{90}Sr) behaves much like calcium and is found in bone tissue (see *The skeleton, an important target*), while caesium-137 (^{137}Cs) follows practically the same metabolic pathways as potassium and so is diffused in muscle tissue.

It is noteworthy that substances that are involved in normal physiological function can nevertheless be toxic if they are present in the body in abnormal quantities – eight or ten litres of water ingested rapidly will induce a coma –, or introduced by a non-physiological route – intravenous injection of pure water induces **lysis** of red cells that can alter renal function.

The toxicity of the exogenous substance is related to its physicochemical properties towards the biological

medium it has entered: carbon monoxide competes with oxygen to bind to haemoglobin with which its chemical bond is stronger; cyanide, by its high chemical affinity for ferric iron, binds to the iron in **mitochondrial** cytochrome oxidase thereby hindering cell respiration; particles of plutonium deposited in the pulmonary alveoli irradiate neighbouring cells; sodium hydroxide crystals in contact with the skin or mucosa will cause burns, as will **beta/gamma emitting radionuclides**.

The toxic effects can vary according to the relative amount of toxin in play, but also the biological parameters of the individual concerned: sex, age, pre-existing health disorders, etc.

Ever since Antiquity, besides preventive action, attempts have been made to forestall or limit the effects of toxins by controlling their entry, diffusion, binding or elimination.

However, once they have entered the body toxins seldom occur in their free forms. Thus using current methods it is fairly usual to encounter pharmacological difficulties mobilising them, whence the importance of intervening promptly at the entry route.

There are various means to do this: mechanical action, chemical neutralisation, administration of a non-toxic substance that competes metabolically with the exogenous element, use of **chelating agents**, i.e., chemical entities able to associate strongly with the toxin to yield a form that is stable in biological media and easily eliminated in urine, for example.

Treatment at several levels

Therapy can be applied at several levels, from the entry of the toxin to its elimination, via its diffusion and binding.

Entry

Neutralisation, as calcium fluoride, of splashes of hydrofluoric acid on the skin using calcium gluconate; removal of **radioactive** particles from the skin using appropriate **surfactants**; inhalation of a DTPA (diethylene triamine penta-acetic acid) **aerosol** to chelate plutonium; precipitation of ^{137}Cs in the intestinal lumen by ingestion of **Prussian blue** (drug name: Radiogardase®).

In highly exceptional circumstances involving pulmonary deposition of toxic materials, broncho-alveolar washing has been indicated.

A particular case is that of the administration of stable iodine, which by rapidly saturating the thyroid gland and temporarily blocking its function, prevents it from subsequently absorbing radioactive iodine.

Diffusion

Generally, the bloodstream is the preferred site of action of chelating agents: the injection of $\text{EDTA Na}_2\text{Ca}$ (an ethylene diamine tetra-acetate), which chelates lead, or of DTPA, which chelates plutonium, causes the elimination of the metal in urine.

Binding

Other than treatments that displace the cyanide ion, or hyperbaric therapy for carbon monoxide poisoning, there are few methods if any that are really effective at this level.



Cyrille Dupont/CEA

Part of the therapeutic arsenal currently available to respond to contamination by nuclear toxics. It includes, in particular, Prussian blue for the chemical neutralisation of certain radionuclides, chelating agents based on DTPA to trap plutonium travelling through the body and facilitate its elimination, and potassium iodide to saturate the thyroid gland with non-radioactive iodine. In this field, researchers are working on new chemical entities able to bind even more efficiently to toxics to form complexes stable in biological media.




Cyrille Dupont/CEA

DTPA (diethylene triamine penta-acetic acid), a plutonium chelating agent that facilitates excretion in urine, can be inhaled in an aerosol or administered by the intravenous route.



Cyrille Dupont/CEA

Bottle of gelatine capsules containing Prussian blue, which when ingested precipitates caesium-137 and thallium-204 in the intestinal lumen and prevents their absorption. It is an iron ferrocyanide better known as a paint pigment.



Elimination

The action most often consists in increasing the natural level of urinary or faecal elimination: for example increasing water intake to clear **incorporated** tritium. Diuretics or laxatives have been used in some cases.

Efficacious but safe

The treatments used in toxicology, like any therapy, need to be efficacious but not harmful. Their own toxicity can be a problem. For example the affinity of chelating agents is seldom exclusive: their use can result in unwanted depletions of vital elements. Harm can also arise from excessive efficacy of the desired effect:

a too-massive **decorporation** of lead may saturate renal tissue and damage the kidney. For this reason special formulations of DTPA have, for example, been developed to avert unwanted calcium or zinc depletion. The application of these treatments must therefore always be preceded by a risk-benefit evaluation.

> Dr **Jean-Michel Giraud**
Medical Adviser

Human Resources and Social Relations Division

> Dr **Jean Piechowski**
Cabinet of the High Commissioner
for Atomic Energy
CEA Headquarters (Paris)

A Natural and artificial radioactivity

Everything on the earth's surface has always been exposed to the action of **ionising radiation** from natural sources. **Natural radiation**, which accounts for 85.5% of total radioactivity (natural plus artificial), is made up of 71% **telluric radiation** and about 14.5% **cosmic radiation**. The **radionuclides** formed by the interaction of **cosmic rays** arriving from stars, and especially the Sun, with the nuclei of elements present in the atmosphere (oxygen and nitrogen) are, in decreasing order of **dose** (Box F, *From rays to dose*) received by the population, carbon-14, beryllium-7, sodium-22 and tritium (hydrogen-3). The last two are responsible for only very low doses.

Carbon-14, with a **half life** of 5,730 years, is found in the human body. Its **activity** per unit mass of carbon has varied over time: it has diminished as carbon dioxide emissions from the combustion of fossil fuels have risen, then was increased by atmospheric nuclear weapon tests.

Beryllium-7, with a half life of 53.6 days, falls onto the leaf surfaces of plants and enters the body by **ingestion** (Box B, *Human exposure routes*). About 50 Bq (becquerels) per person per year of beryllium-7 are ingested.

The main or "primordial" radionuclides are potassium-40, uranium-238 and thorium-232. Along with their radioactive decay products, these elements are present in rocks and soil and are therefore found in many building materials. Their concentrations are generally very low, but vary according to the nature of the mineral. The **gamma radiation** emitted by these radionuclides forms the **telluric radiation**, which is responsible for the **external exposure** of the body. The primordial radionuclides and many of their long-lived descendants

are also found in trace amounts in drinking water and plants: this results in an **internal exposure** by ingestion, plus an additional low exposure by **inhalation** of airborne suspended dust particles.

Potassium-40 is a **beta** and **gamma** emitter with a half life of 1.2 thousand million years, and has no radioactive descendants. This radioactive **isotope** makes up 0.0118% of all natural potassium, and enters the body by ingestion. The mass of natural potassium in the human body is independent of the quantity ingested.

Uranium-238 is an **alpha** emitter with a half life of 4.47 thousand million years. It has thirteen main alpha-, beta- and gamma-emitting radioactive descendants, including **radon-222** (3.82 days) and **uranium-234** (0.246 million years). Uranium-238 and its two descendants **thorium-234** (24.1 days) and **protactinium-234m**⁽¹⁾ (1.18 min), and **uranium-234** are essentially incorporated by ingestion and are mainly concentrated in the bones and kidneys. **Thorium-230**, derived from uranium-234, is an alpha emitter with a period of 80,000 years. It is an **osteotrope**, but enters the body mainly by the pulmonary route (inhalation). **Radium-226**, a descendant of thorium-230, is an alpha emitter with a half life of 1,600 years. It is also an osteotrope and enters the body mainly *via* food. Another osteotrope, **lead-210** (22.3 years), is incorporated by inhalation though mostly by ingestion.

Thorium-232 is an alpha emitter with a half life of 14.1 thousand million

years. It possesses ten main alpha-, beta- and gamma-emitting radioactive descendants including **radon-220** (55 s). Thorium-232 enters the body mainly by inhalation. **Radium-228**, a direct descendant of thorium-232, is a beta-emitter with a half life of 5.75 years. It enters the body mainly in food.

Radon, a gaseous radioactive descendant of uranium-238 and thorium-232, emanates from the soil and building materials, and along with its short-lived alpha-emitting descendants constitutes a source of internal exposure through inhalation. Radon is the most abundant source of natural radiation (about 40% of total radioactivity).

The human body contains nearly 4,500 Bq of potassium-40, 3,700 Bq of carbon-14 and 13 Bq of radium-226 essentially imported in food.

Natural radiation is supplemented by an **anthropic component**, resulting from the medical applications of ionising radiation and to a lesser extent from the nuclear industry. It accounts for about 14.5% of the total radioactivity worldwide, but much more in the developed countries. In the medical field (more than 1 mSv/year on average in France), irradiation by external sources predominates: radiodiagnosis (X-rays) and radiotherapy, long based on caesium-137 and cobalt-60 sources, but now more and more often using linear accelerators. Irradiation by internal routes (curietherapy with iridium-192) has more specialised indications (cervical cancer, for example). The metabolic and physico-chemical properties of some twenty radionuclides are put to use for **medical activities** and in **biological research**. The medical applications comprise radiodiagnostics (**scintigraphy** and radio-

(1) m for metastable. A nuclide is said metastable when a transition delay exists between the excited state of the atom and the stable one.

immunology), and treatment, including thyroid disorders using iodine-131, radioimmunotherapy in certain blood diseases (phosphorus-32) and the treatment of bone metastasis with strontium-89 or radiolabelled phosphonates alongside other uses of radiopharmaceuticals. Among the most widely used radionuclides are: **technetium-99m** (half life 6.02 hours) and **thallium-201** (half life 3.04 days) (scintigraphy), **iodine-131** (half life 8.04 days) (treatment of hyperthyroidism), **iodine-125** (half life 60.14 days) (radioimmunology), **cobalt-60** (half life 5.27 years) (radiotherapy), and **iridium-192** (half life 73.82 days) (curietherapy). The average contribution of radiological examinations to total radioactivity amounts to 14.2%.

The **early atmospheric nuclear weapon tests** scattered fallout over the whole of the earth's surface and caused the exposure of populations and the **contamination** of the food chain by a certain number of radionuclides, most of which, given their short radioactive half lives, have now vanished. There remain **cæsius-137** (30 years), **strontium-90** (29.12 years), some **krypton-85** (10.4 years) and **tritium** (12.35 years), and the isotopes of **plutonium** (half lives 87.7 years to 24,100 years). Currently, the doses corresponding to the fallout from these tests are essentially attributable to **fission products** (cæsius-137) and to carbon-14, rather than **activation products** and plutonium.

In the **Chernobyl accident** (Ukraine), which occurred in 1986, the total radioactivity dispersed into the atmosphere was of the order of 12 milliard milliard (10^{18}) becquerels over a period of 10 days. Three categories of radionu-

clides were disseminated. The first consisted of volatile fission products such as **iodine-131**, **iodine-133** (20.8 hours), **cæsius-134** (2.06 years), **cæsius-137**, **tellurium-132** (3.26 days). The second was composed of solid fission products and **actinides** released in much smaller amounts, in particular the strontium isotopes ^{89}Sr (half life 50.5 days) and ^{90}Sr , the ruthenium isotopes ^{103}Ru (half life 39.3 days) and ^{106}Ru (half life 368.2 days), and **plutonium-239** (24,100 years). The third category was rare gases which although they represented most of the activity released, were rapidly diluted in the atmosphere. They were mainly **xenon-133** (5.24 days) and **krypton-85**.

The contributions of the early atmospheric nuclear weapon tests and the Chernobyl accident to the total radioactivity are roughly 0.2% (0.005 mSv) and 0.07% (0.002 mSv) respectively.

The whole of the **nuclear-powered electricity production** cycle represents only about 0.007% of total radioactivity. Almost all the radionuclides remain confined inside the nuclear reactors and the **fuel** cycle plants. In a nuclear reactor, the reactions that take place inside the fuel yield **transuranics**. **Uranium-238**, which is non-**fissile**, can capture neutrons to give in particular plutonium isotopes ^{239}Pu , ^{240}Pu (half life 6,560 years) and ^{241}Pu (half life 14.4 years), and **americium-241** (432.7 years). The main fission products generated by the fission of **uranium-235** (704 million years) and **plutonium-239** are **iodine-131**, **cæsius-134**, **cæsius-137**, **strontium-90** and **selenium-79** (1.1 million years).

The main radionuclides present in releases, which are performed in a



Laurence Médard/CEA

Classical scintigraphy performed at the Frédéric-Joliot Hospital Service (SHFJ). The gamma-ray camera is used for functional imaging of an organ after administration, usually by the intravenous route, of a radioactive drug (radiopharmaceutical) to the patient. The radionuclides used are specific to the organ being studied: for example, technetium-99m for the kidneys and bones, thallium-201 for the myocardium. The injected radiopharmaceutical emits gamma photons, which are captured by two planar detectors placed at 180° or 45° according to the examination.

very strict regulatory framework are, in liquid release, **tritium**, **cobalt-58** (70.8 days), **cobalt-60**, **iodine-131**, **cæsius-134**, **cæsius-137** and **silver-110m** (249.9 days). In gaseous releases **carbon-14** is the most abundant radionuclide, emitted most often as carbon dioxide. In all the reactors in the world, the total production of radiocarbon dioxide amounts to one tenth of the annual production formed naturally by cosmic radiation.

In addition, certain radionuclides related to the nuclear industry exhibit **chemical toxicity** (Box D, **Radiological and chemical toxicity**).

B Human exposure routes

Human **exposure**, i.e., the effect on the body of a chemical, physical or radiological agent (irrespective of whether there is actual contact), can be external or internal. In the case of **ionising radiation**, exposure results in an energy input to all or part of the body. There can be direct **external irradiation** when the subject is in the path of radiation emitted by a radioactive source located outside the body. The person can be irradiated directly or after reflection off nearby surfaces.

The irradiation can be **acute** or **chronic**. The term **contamination** is used to designate the deposition of matter (here **radioactive**) on structures, surfaces, objects or, as here, a living organism. Radiological contamination, attributable to the presence of **radionuclides**, can occur by the **external** route from the

receptor medium (air, water) and vector media (soils, sediments, plant cover, materials) by contact with skin and hair (cutaneous contamination), or by the **internal** route when the radionuclides are **intaken**, by **inhalation** (gas, particles) from the atmosphere, by **ingestion**, mainly from foods and beverages (water, milk), or by penetration (injury, burns or diffusion through the skin). The term **intoxication** is used when the toxicity in question is essentially chemical.

In the case of **internal contamination**, the dose delivered to the body over time (called the **committed dose**) is calculated for 50 years in adults, and until age 70 years in children. The parameters taken into account for the calculation are: the nature and the intaken quantity of the radionuclide (RN), its

chemical form, its **effective half life**⁽¹⁾ in the body (combination of **physical** and **biological half lives**), the type of **radiation**, the mode of exposure (inhalation, ingestion, injury, transcutaneous), the distribution in the body (deposition in target organs or even distribution), the radiosensitivity of the tissues and the age of the contaminated subject.

Lastly, the **radiotoxicity** is the toxicity due to the ionising radiation emitted by the inhaled or ingested radionuclide. The misleading variable called **potential radiotoxicity** is a *radiotoxic inventory* that is difficult to evaluate and made imprecise by many uncertainties.

(1) The effective half life (T_e) is calculated from the physical half life (T_p) and the biological half life (T_b) by $1 / T_e = 1 / T_p + 1 / T_b$.

F From rays to dose

Radioactivity is a process by which certain naturally-occurring or artificial **nuclides** (in particular those created by **fission**, the splitting of a heavy nucleus into two smaller ones) undergo spontaneous **decay**, with a release of energy, generally resulting in the formation of new nuclides. Termed **radionuclides** for this reason, they are unstable owing to the number of nucleons they contain (protons and neutrons) or their energy state. This decay process is accompanied by the emission of one or more types of **radiation**, ionising or non-ionising, and (or) particles. **Ionising radiation** is electromagnetic or corpuscular radiation that has sufficient energy to ionise certain atoms of the matter in its path by stripping electrons from them. This process can be *direct* (the case with alpha particles) or *indirect* (gamma rays and neutrons).

Alpha radiation, consisting of helium-4 nuclei (two protons and two neutrons), has low penetrating power and is stopped by a sheet of paper or the outermost layers of the skin. Its path in biological tissues is no longer than a few tens of micrometres. This radiation is therefore strongly ionising, i.e., it easily strips electrons from the atoms in the matter it travels through, because the particles shed all their energy over a short distance. For this reason, the hazard due to

radionuclides that are **alpha emitters** is **internal exposure**.

Beta radiation, made up of electrons (beta minus radioactivity) or positrons (beta plus radioactivity), has moderate penetrating power. The particles emitted by **beta emitters** are stopped by a few metres of air, aluminium foil, or a few millimetres of biological tissue. They can therefore penetrate the outer layers of the skin.

Gamma radiation composed of high energy photons, which are weakly ionising but have high penetrating power (more than the **X-ray** photons used in radiodiagnosis), can travel through hundreds of meters of air. Thick shielding of concrete or lead is necessary to protect persons.

The interaction of **neutron radiation** is random, and so it is stopped only by a considerable thickness of concrete, water or paraffin wax. As it is electrically neutral, a neutron is stopped in air by the nuclei of light elements, the mass of which is close to that of the neutron.

- The quantity of energy delivered by radiation is the **dose**, which is evaluated in different ways, according to whether it takes into account the quantity of energy absorbed, its rate of delivery, or its biological effects.

- The **absorbed dose** is the quantity of energy absorbed at a point per unit mass of matter (inert or living),

according to the definition of the International Commission on Radiation Units and Measurements (**ICRU**). It is expressed in **grays** (Gy): 1 gray is equal to an absorbed energy of 1 joule per kilogramme of matter. The *organ absorbed dose* is obtained by averaging the doses absorbed at different points according to the definition of the International Commission on Radiological Protection (**ICRP**).

- The **dose rate**, dose divided by time, measures the intensity of the irradiation (energy absorbed by the matter per unit mass and per unit time). The legal unit is the gray per second (Gy/s), but the gray per minute (Gy/min) is commonly used. Also, radiation has a higher **relative biological effectiveness (RBE)** if the effects produced by the same dose are greater or when the dose necessary to produce a given effect is lower.

- The **dose equivalent** is equal to the dose absorbed in a tissue or organ multiplied by a **weighting factor**, which differs according to the nature of the radiation energy, and which ranges from 1 to 20. Alpha radiation is considered to be 20 times more harmful than gamma radiation in terms of its biological efficiency in producing random (or **stochastic**) effects. The equivalent dose is expressed in sieverts (Sv).

- The **effective dose** is a quantity introduced to try to evaluate harm

F (next)



Foulon/CEA

Technicians operating remote handling equipment on a line at the Atalante facility at CEA Marcoule. The shielding of the lines stops radiation. The operators wear personal dosimeters to monitor the efficacy of the protection.

in terms of whole-body stochastic effects. It is the sum of *equivalent doses* received by the different organs and tissues of an individual, weighted by a factor specific to each of them (weighting factors) according to its specific sensitivity. It makes it possible to sum doses from different sources, and both external and internal radiation. For internal exposure situations (*inhalation, ingestion*), the effective dose is calculated on the basis of the number of *becquerels*

incorporated of a given radionuclide (**DPUI, dose per unit intake**). It is expressed in sieverts (Sv).

- The **committed dose**, as a result of internal exposure, is the cumulated dose received in fifty years (for workers and adults) or until age 70 (for those aged below 20) after the year of **incorporation** of the radionuclide, unless it has disappeared by physical shedding or biological elimination.
- The **collective dose** is the dose received by a population, defined

as the product of the number of individuals (e.g., those working in a nuclear plant, where it is a useful parameter in the optimisation and application of the ALARA system) and the average equivalent or effective dose received by that population, or as the sum of the individual effective doses received. It is expressed in man-sieverts (man.Sv). It should be used only for groups that are relatively homogeneous as regards the nature of their exposure.

D Radiological and chemical toxicity

The chemical toxics linked to the nuclear industry include **uranium** (U), **cobalt** (Co), **boron** (B), used for its neutron-absorbing properties in the heat-exchange fluids of nuclear power plants, **beryllium** (Be), used to slow neutrons, and **cadmium** (Cd), used to capture them. Boron is essential for the growth of plants. Cadmium, like lead (Pb), produces toxic effects on the central nervous system. When the toxicity of an element can be both radiological and chemical, for example that of plutonium (Pu), uranium, neptunium, technetium or cobalt, it is necessary whenever possible to determine what toxic effects are radiological, what are chemical, and what can be either radiological or chemical (see *Limits of the comparison between radiological and chemical hazards*).

For **radioactive** elements with long physical **half lives**, the chemical toxicity is a much greater hazard than the radiological toxicity, as exemplified by rubidium (Rb) and natural uranium.

Thus the chemical toxicity of uranium, which is more important than its radiological toxicity, has led the French regulators to set the **ingested** and **inhaled** mass limits for uranium in chemical compounds at 150 mg and 2.5 mg per day respectively, regardless of the **isotopic** composition of the element.

Certain metals or **metalloids** that are non-toxic at low concentrations can become toxic at high concentrations or in their radioactive form. This is the case for cobalt, which can be **genotoxic**, selenium (Se) (naturally incorporated in **proteins** or **RNA**), technetium (Tc) and iodine (I).



Two-dimensional gel electrophoresis image analysis carried out in the course of nuclear toxicology work at CEA Marcoule Centre in the Rhone Valley.