






D Nuclear fusion reactions

To achieve the **fusion** of two light atom **nuclei**, they must be brought in close proximity to each other, whereas they naturally repel one another, since they both bear a positive electric charge. If the energy this fusion releases is to be recovered, the required energy must first be supplied, to break through this barrier, and allow each nucleus to reach the region, very close to the other nucleus, where the nuclear forces make themselves felt, that are able to overcome this electrostatic repulsion, or **Coulomb barrier**. Once this outcome is achieved, the most energetic reaction are those yielding the fused nucleus with the highest binding energy. In the event at hand, this is the case for helium **isotope** ^4H , which comprises four **nucleons** (two **protons** p and two **neutrons** n).

Of the fusion reactions that yield energy as well as generating, on the one hand, a new – **helium** (He) or **tritium** (T) – nucleus and, on the other hand, a nucleon, four are, at first blush, of particular interest.* The first two are attractive, since they only involve **deuterium** (D), the most abundant **hydrogen** isotope on Earth. Mastering them may be the ultimate goal for controlled fusion, however they are by far the hardest to bring about.





The two subsequent ones, yielding the very stable helium-4 nucleus, are particularly energetic:



The fusion reaction that is most easily achieved, exhibiting as it does the highest cross-section, is that involving a deuterium (D) nucleus and a tritium (T) nucleus, their fusion yielding a helium nucleus and a neutron, with respective energies of 3.5 MeV and 14.1 MeV. Thus it is on this reaction, the so-called D–T reaction, that research work on controlled fusion has focused, whether in the context of **inertial-confinement fusion** or of **magnetic-confinement fusion**.

Production of the tritium required is achieved through a fifth reaction, involving lithium and... the neutrons from the D–T reaction.



The primary fuels, i.e. the true raw materials for a reactor, are thus deuterium and lithium (Li).

* These reactions are known as **thermonuclear** reactions since only a temperature of the order of about a hundred million degrees, together with other density and confinement time conditions (see main article), make it possible to bring them about. See page 8 for the table of the main nuclear reactions occurring inside the Sun.

c Elementary particles and fundamental interactions

Neutrinos are the stealthiest particles in the **standard model of particle physics**, the theoretical framework describing all known **elementary particles** and the **fundamental interactions** they mediate (see Table).

The basic constituents of matter, **fermions**, are partitioned into two main categories:

leptons, which do not respond to **strong interaction**, and **quarks**, which are subject to all of the interactions. The six quarks form three pairs (up/down, charmed/strange, beauty/top). In the lepton category, the **charged leptons** (**electron** e^- , **muon** μ , **tau** τ) are involved in the **electromagnetic interaction** and the

weak interaction, while **neutral leptons** (**electron neutrino** ν_e , **muon neutrino** ν_μ , **tau neutrino** ν_τ) are only subject to weak interaction. In the standard model, neutrinos have zero mass, however experiments have shown they do have some mass, though very small, the exact value of which is as yet unknown. Involvement

atom nucleus electron proton charge + 1 masse : 938.272 MeV/c² neutron zero charge mass : 939.565 MeV/c² nucleon quarks					
		leptons able to move freely		quarks held captive in larger particles, they are not observed separately	
Fermions Common matter is made up of particles from this group. Most of these particles were around just after the Big Bang. Presently only to be found in cosmic rays, and around accelerators.	first family	electron Responsible for electricity and chemical reactions. Charge: - 1. mass: 0.511 MeV/c²	electron neutrino Has no electric charge, and interacts very seldom with the ambient medium.	down Electric charge: - 1/3. The proton holds one, the neutron two. mass: 5 - 8,5 MeV/c²	up Electric charge: + 2/3. The proton holds two, the neutron one. mass: 1,5 - 4,5 MeV/c²
	second family	muon A more massive companion to the electron. mass: 105,658 MeV/c²	muon neutrino Properties similar to those of the electron neutrino.	strange A heavier companion to "up." mass: 80 - 155 MeV/c²	charmed A heavier companion to "down." mass: 1 000 - 1 400 MeV/c²
	third family	tau Heavier still. mass: 1 777 MeV/c²	tau neutrino Properties similar to those of the electron neutrino.	beauty Heavier still. mass: 4 000 - 4 500 MeV/c²	top Heaviest in the family (observed in 1985): 180 times the proton mass. mass: 174 300 ± 5 100 MeV/c²
Vector Bosons Fundamental particles carrying out transmission of all natural forces.		photon Elementary grain of light, vector for the electromagnetic force.	gluon Bearer of the strong force between quarks.	W[±], Z⁰ Bearers of the weak force, responsible for some forms of radioactive decay.	
Higgs Boson?		Responsible for "electroweak symmetry breaking."			

Tableau.
Constituants élémentaires.

of the various elementary constituents in the fundamental interactions is governed by their quantum numbers, or interaction charges (electric charge, color charge ⁽¹⁾...). To every constituent of matter is associated its **antiparticle**, a particle having the same mass and opposite charges. The **gravitational force**, which is not included in the standard model, acts on all fermions in proportion to their mass. The table of elementary constituents of matter manifests another classification – independently from their involvement in fundamental interactions – into three generations, or families. From one family to the next, charged quarks and leptons having the same charges only differ by their mass. The electron, up quark and down quark, which all belong to the first family, are the lightest massive particles. They are stable particles, and the constituents of common matter. For instance, the **proton** is made up of two up quarks and one down quark; the **neutron**, of two down quarks and one up quark. Particles in the other two families are unstable, and rapidly decay into

stable first-generation particles. This is why all the stable matter in the Universe is made up from constituents from the first family.

According to quantum mechanics, for an interaction to take place, at least one elementary particle, a **boson**, must be emitted, absorbed or exchanged. The **photon** is the vector for the electromagnetic interaction, the **W⁺**, **W⁻** and **Z⁰** mediate the weak interaction, and **gluons** act as messengers for the strong interaction. Quarks and charged leptons exchange photons, but conserve their electric charge after the exchange, the photon having no electric charge. Since the photon's mass is zero, the electromagnetic interaction's range is infinite. Having no electric charge, neutrinos are the only elementary fermions that are not subject to electromagnetic interaction.

In the electroweak theory (a unification of the weak and electromagnetic interactions), the weak interaction has two aspects: charged-current weak interaction, for which the interaction vectors are the **W⁺** and **W⁻**; and neutral-current weak interaction, for which the mediator is **Z⁰**. These two forms of weak interaction are active between all elementary fermions (quarks, charged leptons and neutrinos). The mass of these bosons being very large (80,000 MeV/c² for **W[±]**, 91,180 MeV/c² for **Z⁰**), the range of the weak interaction is tiny – of the order of

10⁻¹⁸ m. Since **W[±]** bosons have a non-zero electric charge, fermions exchanging such bosons undergo a change in electric charge, as of nature (flavor). Conversely, since the **Z⁰** boson has no electric charge, fermions exchanging one undergo no change in nature. In effect, neutral-current weak interaction is somewhat akin to exchanging a photon. As a general rule, if two fermions are able to exchange a photon, they can also exchange a **Z⁰**. On the other hand, a neutrino has the ability to exchange a **Z⁰** with another particle, though not a photon. Only those quarks that have a color charge exchange gluons, these in turn being bearers of a color charge. Thus, when a gluon exchange takes place between quarks, the latter exchange their respective colors. Gluons have zero mass, however, since they do bear a color charge, they are able to interact. The range of the strong interaction is consequently very restricted – of the order of 10⁻¹⁵ m.

The graviton, the vector for gravitational interaction, has not so far been observed.

Theory predicts that another fundamental interaction mechanism exists, responsible for the mass of elementary particles, for which the messenger is the Higgs boson, which remains as yet undiscovered. This boson makes it possible to assign a mass to elementary fermions of zero mass that interact with it.

(1) Color charge: a quantum number that determines whether a particle is involved in strong interaction. The color charge can take on three values: “red,” “green,” or “blue” – such colors bearing no relation to visible colors. Every quark bears one of the three color charges, every antiquark one of the three anticolor charges. Gluons bear double color–anticolor charges (eight possible combinations).

fundamental interaction	messenger	actions
gravitational	graviton?	responsible for the mutual attraction of any two masses and for the law of falling bodies
electromagnetic	photon	responsible for the attraction between electrons and atomic nuclei, hence for the cohesion of atoms and molecules
weak	W ⁺ , W ⁻ , Z ⁰	the root cause of thermonuclear fusion inside the Sun, ensuring its longevity. β ⁻ and β ⁺ radioactivity, and reactions involving neutrinos are weak interactions
strong	gluons	ensures the cohesion of the atomic nucleus

Table.
Fundamental interaction and elementary constituents.