


F How is solar energy turned into electricity?

The **photovoltaic effect** used in **solar cells** (which are further assembled into **modules**, then into **panels**) for the direct conversion of the rays of sunlight into electricity involves the generation and transport of positive and negative electric charges through the action of light in a **semiconducting** material.

Silicon was selected for its electronic properties, characterized by presence of four **electrons** in its outermost shell (column IV of the Mendeleyev periodic table). In solid silicon, each **atom** is bound to four neighbors, and all the electrons in the outermost shell participate in the bonds. If a silicon atom is substituted for by an atom from column V (a phosphorus atom, for instance), one of the electrons is not involved in the bonds; it is thus free to move through the lattice. There is electron conduction, and the semiconductor is designated as an *n-type doped*

semiconductor. If, on the other hand, a **silicon** atom is substituted for by an atom from column III (boron, for instance), one electron is missing if all bonds are to be maintained, and an electron can move in to fill this gap. One then speaks of hole conduction, and the semiconductor is said to be a *p-type doped semiconductor*. Atoms of elements such as boron or phosphorus are **doping agents** in silicon.



When an *n*-type semiconductor is brought into contact with a *p*-type semiconductor, the excess electrons from the *n* material diffuse into the *p* material. The initially *n*-doped region becomes positively charged, while the initially *p*-doped region takes on a negative charge. An electric field is thus set up between the *n* and *p* regions, tending to force electrons back into the *n* region, and equilibrium arises. A **junction** has been set up, and, if metallic contacts are placed onto the *n* and *p* regions,

a **diode** is obtained.

When this diode is illuminated, **photons** are absorbed by the material, and every photon gives rise to an electron and a hole (this is termed an **electron-hole pair**). The diode junction keeps apart the electrons and the holes, giving rise to a potential difference between the *n* and *p* contacts, and a current passes, if a resistance is inserted between the diode contacts (see Figure).

c Elementary particles and fundamental interactions

Neutrinos are the stealthiest particles in the **standard model of particle physics**, the theoretical framework describing all known **elementary particles** and the **fundamental interactions** they mediate (see Table).

The basic constituents of matter, **fermions**, are partitioned into two main categories:

leptons, which do not respond to **strong interaction**, and **quarks**, which are subject to all of the interactions. The six quarks form three pairs (up/down, charmed/strange, beauty/top). In the lepton category, the **charged leptons** (**electron** e^- , **muon** μ , **tau** τ) are involved in the **electromagnetic interaction** and the

weak interaction, while **neutral leptons** (**electron neutrino** ν_e , **muon neutrino** ν_μ , **tau neutrino** ν_τ) are only subject to weak interaction. In the standard model, neutrinos have zero mass, however experiments have shown they do have some mass, though very small, the exact value of which is as yet unknown. Involvement


					
		leptons able to move freely		quarks held captive in larger particles, they are not observed separately	
Fermions Common matter is made up of particles from this group. Most of these particles were around just after the Big Bang. Presently only to be found in cosmic rays, and around accelerators.	first family	electron  Responsible for electricity and chemical reactions. Charge: -1 . mass: $0.511 \text{ MeV}/c^2$	electron neutrino  Has no electric charge, and interacts very seldom with the ambient medium.	down  Electric charge: $-1/3$. The proton holds one, the neutron two. mass: $5 - 8,5 \text{ MeV}/c^2$	up  Electric charge: $+2/3$. The proton holds two, the neutron one. mass: $1,5 - 4,5 \text{ MeV}/c^2$
	second family	muon  A more massive companion to the electron. mass: $105,658 \text{ MeV}/c^2$	muon neutrino  Properties similar to those of the electron neutrino.	strange  A heavier companion to "up." mass: $80 - 155 \text{ MeV}/c^2$	charmed  A heavier companion to "down." mass: $1\,000 - 1\,400 \text{ MeV}/c^2$
	third family	tau  Heavier still. mass: $1\,777 \text{ MeV}/c^2$	tau neutrino  Properties similar to those of the electron neutrino.	beauty  Heavier still. mass: $4\,000 - 4\,500 \text{ MeV}/c^2$	top  Heaviest in the family (observed in 1985): 180 times the proton mass. mass: $174\,300 \pm 5\,100 \text{ MeV}/c^2$
Vector Bosons Fundamental particles carrying out transmission of all natural forces.		photon  Elementary grain of light, vector for the electromagnetic force.	gluon  Bearer of the strong force between quarks.	$W^\pm, Z^0$  Bearers of the weak force, responsible for some forms of radioactive decay.	
Higgs Boson? 		Responsible for "electroweak symmetry breaking."			

Tableau.
Constituants élémentaires.

of the various elementary constituents in the fundamental interactions is governed by their quantum numbers, or interaction charges (electric charge, color charge ⁽¹⁾...). To every constituent of matter is associated its **antiparticle**, a particle having the same mass and opposite charges. The **gravitational force**, which is not included in the standard model, acts on all fermions in proportion to their mass. The table of elementary constituents of matter manifests another classification – independently from their involvement in fundamental interactions – into three generations, or families. From one family to the next, charged quarks and leptons having the same charges only differ by their mass. The electron, up quark and down quark, which all belong to the first family, are the lightest massive particles. They are stable particles, and the constituents of common matter. For instance, the **proton** is made up of two up quarks and one down quark; the **neutron**, of two down quarks and one up quark. Particles in the other two families are unstable, and rapidly decay into

stable first-generation particles. This is why all the stable matter in the Universe is made up from constituents from the first family.

According to quantum mechanics, for an interaction to take place, at least one elementary particle, a **boson**, must be emitted, absorbed or exchanged. The **photon** is the vector for the electromagnetic interaction, the **W⁺**, **W⁻** and **Z⁰** mediate the weak interaction, and **gluons** act as messengers for the strong interaction. Quarks and charged leptons exchange photons, but conserve their electric charge after the exchange, the photon having no electric charge. Since the photon's mass is zero, the electromagnetic interaction's range is infinite. Having no electric charge, neutrinos are the only elementary fermions that are not subject to electromagnetic interaction.

In the electroweak theory (a unification of the weak and electromagnetic interactions), the weak interaction has two aspects: charged-current weak interaction, for which the interaction vectors are the **W⁺** and **W⁻**; and neutral-current weak interaction, for which the mediator is **Z⁰**. These two forms of weak interaction are active between all elementary fermions (quarks, charged leptons and neutrinos). The mass of these bosons being very large (80,000 MeV/c² for **W[±]**, 91,180 MeV/c² for **Z⁰**), the range of the weak interaction is tiny – of the order of

10⁻¹⁸ m. Since **W[±]** bosons have a non-zero electric charge, fermions exchanging such bosons undergo a change in electric charge, as of nature (flavor). Conversely, since the **Z⁰** boson has no electric charge, fermions exchanging one undergo no change in nature. In effect, neutral-current weak interaction is somewhat akin to exchanging a photon. As a general rule, if two fermions are able to exchange a photon, they can also exchange a **Z⁰**. On the other hand, a neutrino has the ability to exchange a **Z⁰** with another particle, though not a photon. Only those quarks that have a color charge exchange gluons, these in turn being bearers of a color charge. Thus, when a gluon exchange takes place between quarks, the latter exchange their respective colors. Gluons have zero mass, however, since they do bear a color charge, they are able to interact. The range of the strong interaction is consequently very restricted – of the order of 10⁻¹⁵ m.

The graviton, the vector for gravitational interaction, has not so far been observed.

Theory predicts that another fundamental interaction mechanism exists, responsible for the mass of elementary particles, for which the messenger is the Higgs boson, which remains as yet undiscovered. This boson makes it possible to assign a mass to elementary fermions of zero mass that interact with it.

(1) Color charge: a quantum number that determines whether a particle is involved in strong interaction. The color charge can take on three values: “red,” “green,” or “blue” – such colors bearing no relation to visible colors. Every quark bears one of the three color charges, every antiquark one of the three anticolor charges. Gluons bear double color–anticolor charges (eight possible combinations).

fundamental interaction	messenger	actions
gravitational	graviton?	responsible for the mutual attraction of any two masses and for the law of falling bodies
electromagnetic	photon	responsible for the attraction between electrons and atomic nuclei, hence for the cohesion of atoms and molecules
weak	W ⁺ , W ⁻ , Z ⁰	the root cause of thermonuclear fusion inside the Sun, ensuring its longevity. β ⁻ and β ⁺ radioactivity, and reactions involving neutrinos are weak interactions
strong	gluons	ensures the cohesion of the atomic nucleus

Table.
Fundamental interaction and elementary constituents.